



### Anti-GAPDH Antibody

Catalog #	Source	Reactivity	Applications
GP0002	Rabbit	H, M, R, Rb, C, Mk, S, X	WB, IH
<b>Description</b>	Rabbit polyclonal to GAPDH		
<b>Immunogen</b>	Recombinant protein corresponding to human GAPDH.		
<b>Purification</b>	The antibody was purified by affinity chromatography.		
<b>Specificity</b>	Recognizes endogenous levels of GAPDH protein.		
<b>Clonality</b>	Polyclonal		
<b>Form</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		
<b>Dilution</b>	WB (1/2000 - 1/5000), IH (1/100 - 1/200)		
<b>Gene Symbol</b>	GAPDH		
<b>Alternative Names</b>	GAPD; Glyceraldehyde-3-phosphate dehydrogenase; GAPDH; Peptidyl-cysteine S-nitrosylase GAPDH		
<b>Entrez Gene</b>	2597 (Human); 100042025 (Mouse); 24383 (Rat)		
<b>SwissProt</b>	P04406 (Human); P16858 (Mouse); P04797 (Rat)		
<b>Storage/Stability</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.		



Western blot analysis of GAPDH expression in 293T (A), NIH3T3 (B), rat brain (C), sheep muscle (D), rabbit testis (E) whole cell lysates.



Immunohistochemical analysis of GAPDH staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.